

Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Cord Blood Methylmercury and Fetal Growth Outcomes in Baltimore Newborns: Potential Confounding and Effect Modification by Omega-3 Fatty Acids, Selenium, and Sex

Ellen M. Wells, Julie B. Herbstman, Yu Hong Lin, Jeffery Jarrett, Carl P. Verdon, Cynthia Ward, Kathleen L. Caldwell, Joseph R. Hibbeln, Frank R. Witter, Rolf U. Halden, and Lynn R. Goldman

Table of Contents

Table S1: Spearman correlation coefficients (*p*-values) for mercury, selenium and n-3 HUFAs, THREE Study, 2004-2005

Table S2: Regression coefficient (95% confidence intervals) for MeHg, n-3 HUFAs, selenium, and IHg in various multivariable regression models, N=271, THREE Study, 2004-2005

Table S3: Adjusted regression coefficient (95% confidence intervals) for ln(MeHg), among the full cohort as well as stratum-specific estimates, THREE Study, 2004-2005